



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification <sup>6</sup> : <b>C07K 5/00, C07D 403/00, 401/00</b>		A2	(11) International Publication Number: <b>WO 98/27108</b>
			(43) International Publication Date: 25 June 1998 (25.06.98)
(21) International Application Number: PCT/JP97/04243		305 (JP). SHIMA, Ichiro [JP/JP]; 5-25-105, Gosyogaoka, Moriya-cho, Kitasouma-gun, Ibaraki 302-01 (JP). OHNE, Kazuhiko [JP/JP]; 1-16-15-A102, Ninomiya, Tsukuba-shi, Ibaraki 305 (JP). YOSHIHARA, Kousei [JP/JP]; 2-4-38-405, Manabe, Tsuchiura-shi, Ibaraki 300 (JP). OKU, Teruo [JP/JP]; 8-2, Midorigaoka, Tsukuba-shi, Ibaraki 305 (JP).  (74) Agent: TAKASHIMA, Hajime; Yuki Building, 3-9, Hiranomachi 3-chome, Chuo-ku, Osaka-shi, Osaka 541 (JP).  (81) Designated States: AU, CA, CN, HU, IL, JP, KR, MX, US, Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).  Published Without international search report and to be republished upon receipt of that report.	
(22) International Filing Date: 20 November 1997 (20.11.97)			
(30) Priority Data:			
PO 4219	16 December 1996 (16.12.96) AU		
PO 5929	1 April 1997 (01.04.97) AU		
PO 9030	9 September 1997 (09.09.97) AU		
(71) Applicant (for all designated States except US): FUJISAWA PHARMACEUTICAL CO., LTD. [JP/JP]; 4-7, Doshomachi 3-chome, Chuo-ku, Osaka-shi, Osaka 541 (JP).			
(71) Applicant (for US only): YATABE, Yoshiko (heiress of the deceased inventor) [JP/JP]; 4-1-1-421-201, Namiki, Tsukuba-shi, Ibaraki 305 (JP).			
(72) Inventor: YATABE, Takumi (deceased).			
(72) Inventors; and			
(75) Inventors/Applicants (for US only): ITOH, Yoshikuni [JP/JP]; 2-49-12, Himuro-cho, Takatsuki-shi, Osaka 569-11 (JP). INOUE, Takayuki [JP/JP]; 4-15-2-2-201, Matsushiro, Tsukuba-shi, Ibaraki 305 (JP). HAMASHIMA, Hitoshi [JP/JP]; 3-25-4-202, Matsushiro, Tsukuba-shi, Ibaraki			
(54) Title: NEW AMIDE COMPOUNDS			
(57) Abstract			
<p>A compound of formula (I) wherein each symbol is as defined in the specification, and pharmaceutically acceptable salts thereof. The compound (I) of the present invention and pharmaceutically acceptable salts thereof possess a strong inhibitory activity on the production of nitric oxide (NO), and are useful for prevention and/or treatment of NO-mediated diseases such as adult respiratory distress syndrome, cardiovascular ischemia, myocarditis, heart failure, synovitis, shock, diabetes, diabetic nephropathy, diabetic retinopathy, diabetic neuropathy, glomerulonephritis, peptic ulcer, inflammatory bowel disease, cerebral infarction, cerebral ischemia, cerebral hemorrhage, migraine, rheumatoid arthritis, gout, neuritis, postherpetic neuralgia, osteoarthritis, osteoporosis, systemic lupus erythematosus, rejection by organ transplantation, asthma, metastasis, Alzheimer's disease, arthritis, CNS disorders, dermatitis, hepatitis, liver cirrhosis, multiple sclerosis, pancreatitis, atherosclerosis, and the like in human being and animals.</p>			
<div style="text-align: center;"> <math display="block">  \begin{array}{c}  R^2 \\    \\  R^1 - CON - (Y) - \begin{array}{c} N \\ \parallel \\ X \end{array} \begin{array}{c} R^5 \\ \parallel \\ R^4 \end{array}  \end{array}  \quad (I)  </math> </div>			

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

DESCRIPTION  
NEW AMIDE COMPOUNDS

TECHNICAL FIELD

This invention relates to new amide compounds and pharmaceutically acceptable salts thereof which are useful as medicament.

BACKGROUND ART

Some peptide compounds have been known as described, for example, in EP 0 394 989 A2.

DISCLOSURE OF INVENTION

This invention relates to new amide compounds.

One object of this invention is to provide the new and useful amide compounds and pharmaceutically acceptable salts thereof which possess a strong inhibitory activity on the production of nitric oxide (NO).


Another object of this invention is to provide a process for the preparation of the amide compounds and salts thereof.

A further object of this invention is to provide a pharmaceutical composition comprising said amide compound or a pharmaceutically acceptable salt thereof.

Still further object of this invention is to provide a use of said amide compounds or pharmaceutically acceptable salts thereof as a medicament for prophylactic and therapeutic treatment of NO-mediated diseases such as adult respiratory distress syndrome, cardiovascular ischemia, myocarditis, heart failure, synovitis, shock (e.g., septic shock, etc.), diabetes (e.g., insulin-dependent diabetes mellitus, etc.), diabetic nephropathy, diabetic retinopathy, diabetic neuropathy, glomerulonephritis, peptic ulcer, inflammatory bowel disease (e.g., ulcerative colitis, chronic colitis, etc.), cerebral

infarction, cerebral ischemia, cerebral hemorrhage, migraine, rheumatoid arthritis, gout, neuritis, postherpetic neuralgia, osteoarthritis, osteoporosis, systemic lupus erythematosus, rejection by organ transplantation, asthma, metastasis, Alzheimer's disease, arthritis, CNS disorders, dermatitis, hepatitis, liver cirrhosis, multiple sclerosis, pancreatitis, atherosclerosis, and the like in human being and animals.

The object amide compounds of the present invention are novel and can be represented by the following general formula (I)



wherein


R<sup>1</sup> is indolyl which may have a suitable substituent selected from the group consisting of lower alkyl, phenyl, halogen, lower alkoxy, and nitro, benzofuranyl, phenyl which may have one or two suitable substituent(s) selected from the group consisting of amino, acylamino, lower alkylamino, halogen, lower alkoxy and nitro, lower alkyl, quinoxaliny, quinolyl, pyrrolyl, pyrimidinyl having benzofuranyl, benzimidazolyl, benzothienyl, benzothiazolyl, benzoxazolyl, indolinyl, anilino, phenylcarbamoyl or imidazolyl which may have one or two suitable substituent(s) selected from the group consisting of phenyl, lower alkyl and indolyl;

R<sup>2</sup> is hydrogen or phenyl(lower)alkyl;

R<sup>4</sup> is hydrogen, phenyl or pyridyl, each of which may have suitable substituent(s) selected from the group consisting of lower alkyl, lower alkoxy, lower alkylthio, halogen, trihalomethyl, nitro, cyano, imidazolyl, optionally protected hydroxy, acyl, amino, acylamino, diacylamino, di(lower)alkylamino, amino(lower)alkyl, acylamino(lower)alkyl, pyrazolyl,


morpholinyl, piperidyl, triazolyl, lower alkoxy(lower)alkoxy, hydroxy(lower)alkyl, lower alkylpiperazinyl, phenyl and carboxy, quinolyl or 3,4-methylenedioxyphenyl;

R<sup>5</sup> is hydrogen, imidazolyl, phenyl, nitrophenyl, phenyl(lower)alkyl, optionally esterified carboxy or a group of the formula



in which R<sup>7</sup> and R<sup>8</sup> are the same or different and each is hydrogen, phenyl, phenyl(lower)alkyl, lower alkyl or lower alkoxy; or


R<sup>4</sup> and R<sup>5</sup> in combination form a group of the formula



Y is a group of the formula




in which R<sup>3</sup> is hydrogen or a group of the formula



in which R<sup>6</sup> is optionally protected hydroxy, acyl, carboxy, acylamino, lower alkoxy, phenyl(lower)alkoxy, lower alkylthio, phenyl which may have a suitable substituent selected from the group consisting of lower alkoxy, halogen, amino, acylamino, diacylamino and nitro, pyridyl which may have a suitable substituent selected from the group consisting of lower alkoxy and halogen, pyrazinyl, pyrimidinyl, furyl, imidazolyl, naphthyl, N-(lower)-alkylindolyl or 3,4-methylenedioxyphenyl, and n is an integer of 0 to 3,

or a group of the formula



in which R<sup>11</sup> is phenyl, phenoxy or phenyl(lower)alkoxy; or


R<sup>2</sup> and R<sup>3</sup> in combination form a group of the formula




m is 0 or 1; and

X is S or NR<sup>9</sup>

in which R<sup>9</sup> is hydrogen, lower alkyl, cyclo(lower)alkyl or a group of the formula




in which R<sup>10</sup> is hydrogen, lower alkyl or lower alkoxy;  
or a pharmaceutically acceptable salt thereof,  
provided that the compound shown below is excluded:  
a compound of the formula



wherein


R<sup>1'</sup> is indolyl or benzofuranyl;

R<sup>2'</sup> is hydrogen, lower alkylthio(lower)alkyl or a group of the formula



in which R<sup>5'</sup> is hydrogen, lower alkoxy or halogen;  
R<sup>3'</sup> is hydrogen, quinolyl or phenyl which may have a suitable  
substituent selected from the group consisting of lower alkyl,  
lower alkoxy, lower alkylthio and halogen;  
R<sup>4'</sup> is hydrogen or optionally esterified carboxy; and  
X' is S or NR<sup>6'</sup>

in which R<sup>6'</sup> is hydrogen, lower alkyl or a group of the formula



in which R<sup>7'</sup> is lower alkyl or lower alkoxy,  
and a pharmaceutically acceptable salt thereof.

Suitable pharmaceutically acceptable salts of the object compound (I) are conventional non-toxic salts and include, for example, a salt with a base or an acid addition salt such as a salt with an inorganic base, for example, an alkali metal salt (e.g., sodium salt, potassium salt, etc.), an alkaline earth metal salt (e.g., calcium salt, magnesium salt, etc.), an ammonium salt; a salt with an organic base, for example, an organic amine salt (e.g., triethylamine salt, pyridine salt, picoline salt, ethanolamine salt, triethanolamine salt, dicyclohexylamine salt, N,N'-dibenzylethylenediamine salt, etc.); an inorganic acid addition salt (e.g., hydrochloride, hydrobromide, sulfate, phosphate, etc.); an organic carboxylic or sulfonic acid addition salt (e.g., formate, acetate, trifluoroacetate, maleate, tartrate, citrate, fumarate, methanesulfonate, benzenesulfonate, toluenesulfonate, etc.); and a salt with a basic or acidic amino acid (e.g., arginine, aspartic acid, glutamic acid, etc.).

In the above and subsequent descriptions of the present specification, suitable examples and illustration of the various definitions which the present invention intends to include within the scope thereof are explained in detail as follows.

The term "lower" is used to intend a group having 1 to 6, preferably 1 to 4, carbon atom(s), unless otherwise provided.

Suitable "lower alkyl" and "lower alkyl moiety" in the terms "lower alkylthio", "lower alkylthio(lower)alkyl", "N-(lower)-alkylindolyl", "lower alkylamino", "di(lower)alkylamino",

"phenyl(lower)alkyl", "amino(lower)alkyl", "acylamino(lower)alkyl", "hydroxy(lower)alkyl" and "lower alkylpiperazinyl" include straight or branched one having 1 to 6 carbon atom(s), such as methyl, ethyl, propyl, isopropyl, butyl, isobutyl, sec-butyl, tert-butyl, pentyl, tert-pentyl and hexyl, in which more preferred one is C<sub>1</sub>-C<sub>4</sub> alkyl.

Suitable "lower alkoxy" and "lower alkoxy moiety" in the terms "lower alkoxy(lower)alkoxy" and "phenyl(lower)alkoxy" include, for example, methoxy, ethoxy, propoxy, isopropoxy, butoxy, isobutoxy, tert-butoxy, pentyloxy, tert-pentyloxy and hexyloxy, in which more preferred one is C<sub>1</sub>-C<sub>4</sub> alkoxy.

Suitable "halogen" includes, for example, fluorine, bromine, chlorine and iodine.

"Optionally esterified carboxy" includes carboxy and esterified carboxy. Suitable examples of said ester include lower alkyl ester (e.g., methyl ester, ethyl ester, propyl ester, isopropyl ester, butyl ester, isobutyl ester, tert-butyl ester, pentyl ester, tert-pentyl ester, hexyl ester, etc.); lower alkenyl ester (e.g., vinyl ester, allyl ester, etc.); lower alkynyl ester (e.g., ethynyl ester, propynyl ester, etc.); lower alkoxy(lower)alkyl ester (e.g., methoxymethyl ester, ethoxymethyl ester, isopropoxymethyl ester, 1-methoxyethyl ester, 1-ethoxyethyl ester, etc.); mono(or di or tri)-aryl(lower)alkyl ester, for example, mono(or di or tri)phenyl(lower)-alkyl ester which may have one or more suitable substituent(s) [e.g., benzyl ester, 4-methoxybenzyl ester, 4-nitrobenzyl ester, phenethyl ester, trityl ester, benzhydryl ester, bis(methoxyphenyl)methyl ester, 3,4-dimethoxybenzyl ester, 4-hydroxy-3,5-di-tert-butylbenzyl ester, etc.]; and aryl ester which may have one or more suitable substituent(s) such as substituted or unsubstituted phenyl ester (e.g., phenyl ester, tolyl ester, tert-butylphenyl ester, xylyl ester, mesityl ester, cumenyl ester, 4-chlorophenyl ester, 4-methoxyphenyl ester, etc.).

Suitable "trihalomethyl" includes, for example, trifluoromethyl,




trichloromethyl and tribromomethyl, in which preferred one is trifluoromethyl.

Suitable "amino protective group" includes, for example, acyl and conventional protective group such as mono(or di or tri)aryl(lower)-alkyl, for example, mono(or di or tri)phenyl(lower)alkyl (e.g., benzyl, trityl, etc.).

Suitable "acyl" and "acyl moiety" in the terms "acylamino", "diacylamino" and "acylamino(lower)alkyl" include, for example, carbamoyl which may be substituted by suitable substituent(s), aliphatic acyl group and acyl group containing an aromatic ring, which is referred to as aromatic acyl, or a heterocyclic ring, which is referred to as heterocyclic acyl.

Suitable examples of said acyl are illustrated as follows:  
"carbamoyl which may be substituted by suitable substituent(s)" includes a group of the formula



wherein  $\text{R}^{1'2}$  and  $\text{R}^{1'3}$  are the same or different and each is hydrogen, lower alkyl, phenyl which may have a suitable substituent selected from the group consisting of lower alkoxy and halogen, phenyl(lower)-alkyl, pyridyl, pyridyl(lower)alkyl or 3,4-methylenedioxyphenyl; aliphatic acyl such as lower alkanoyl which may be substituted by one to three halogen atoms (e.g., formyl, acetyl, propanoyl, butanoyl, 2-methylpropanoyl, pentanoyl, 2,2-dimethylpropanoyl, hexanoyl, trichloroacetyl, trifluoroacetyl, etc.), lower alkoxycarbonyl (e.g., methoxycarbonyl, ethoxycarbonyl, tert-butoxycarbonyl, tert-pentyloxycarbonyl, etc.), lower alkylsulfonyl (e.g., methylsulfonyl, ethylsulfonyl, etc.), lower alkoxysulfonyl (e.g., methoxysulfonyl, ethoxysulfonyl, etc.), cyclo(lower)alkylcarbonyl (e.g., cyclopentylcarbonyl, cyclohexylcarbonyl, etc.), and the like; aromatic acyl such as aroyl (e.g., benzoyl, toluoyl, naphthoyl, etc.), aryl(lower)alkanoyl [e.g., phenyl(lower)alkanoyl (e.g., phenylacetyl,



phenylpropanoyl, phenylbutanoyl, etc.), naphthyl(lower)alkanoyl (e.g., naphthylacetyl, naphthylpropanoyl, naphthylbutanoyl, etc.), etc.], aryl(lower)alkoxycarbonyl [e.g., phenyl(lower)alkoxycarbonyl (e.g., benzyloxycarbonyl, etc.), etc.], aryloxycarbonyl (e.g., phenoxycarbonyl, naphthyloxycarbonyl, etc.), aryloxy(lower)alkanoyl (e.g., phenoxyacetyl, phenoxypropionyl, etc.), arylsulfonyl (e.g., phenylsulfonyl, p-tolylsulfonyl, etc.), and the like; heterocyclic acyl such as indolylcarbonyl (e.g., indolyl-2-ylcarbonyl, etc.), benzofuranylcarbonyl (e.g., benzofuran-2-ylcarbonyl), quinoxalinyllcarbonyl, quinolylcarbonyl, pyrrolylcarbonyl, benzimidazolylcarbonyl, benzothienylcarbonyl, benzothiazolylcarbonyl, imidazolylcarbonyl, pyridylcarbonyl, morpholinylcarbonyl (e.g., morpholinocarbonyl) and the like.

"Optionally protected hydroxy" includes hydroxy and protected hydroxy. Suitable examples of "hydroxy protective group" in the term "protected hydroxy" include acyl (e.g., acetyl, trichloroacetyl, etc.), mono(or di or tri)phenyl(lower)alkyl which may have one or more suitable substituent(s) (e.g., benzyl, 4-methoxybenzyl, trityl, etc.), trisubstituted silyl [e.g., tri(lower)alkylsilyl (e.g., trimethylsilyl, tert-butyldimethylsilyl, etc.), etc.], tetrahydropyranyl and the like.

Suitable "protected carboxy" is carboxy group protected by conventional protective group such as lower alkoxycarbonyl [e.g., methoxycarbonyl, ethoxycarbonyl, propoxycarbonyl, isopropoxycarbonyl, butoxycarbonyl, sec-butoxycarbonyl, isobutoxycarbonyl, tert-butoxycarbonyl, pentyloxycarbonyl, neopentyloxycarbonyl, hexyloxycarbonyl, etc.], optionally substituted phenyl(lower)-alkoxycarbonyl for example, mono- or di- or triphenyl(lower)-alkoxycarbonyl which may be substituted by nitro [e.g., benzyloxycarbonyl, 4-nitrobenzyloxycarbonyl, benzhydryloxycarbonyl, trityloxycarbonyl, etc.] and the like.

Suitable "cyclo(lower)alkyl" includes cycloalkyl having 3 to 6

Table

Example No.	Formula
27	
	
28	